

# Hepatitis C Enrollment Form

Ph: 806-324-5447 • Toll Free 866-629-6779  
 Toll Free Fax: 866-217-8034  
 Email: [specialty@maxor.com](mailto:specialty@maxor.com)  
[www.maxorspecialty.com](http://www.maxorspecialty.com)



| PATIENT INFORMATION   |  | PRESCRIBER INFORMATION   |               |         |
|---|--|--|---------------|---------|
| Patient Name _____<br>Date of Birth _____ <input type="radio"/> Male <input type="radio"/> Female<br>Street Address _____ Apt # _____<br>City _____ State _____ Zip _____<br>Phone-Primary _____ Secondary _____<br><input type="radio"/> English <input type="radio"/> Spanish <input type="radio"/> Other _____<br>Email Address _____<br>Social Security # _____<br><input type="radio"/> NKDA Known Drug Allergies _____<br>Weight _____ kg/lb Height _____ in/cm<br>Please attach front and back of patient's insurance cards  |  | Physician Name _____<br>NPI _____ License # _____<br>Office Contact _____<br>Street Address _____ Ste # _____<br>City _____ State _____ Zip _____<br>Phone _____ Fax _____<br>By signing this form, I authorize Maxor Specialty to act as my agent for prior authorizations & prescription reimbursement for the above listed patient.<br>Physician Signature _____<br>Date _____<br><input type="radio"/> Dispense as written <input type="radio"/> Product substitution permitted<br>** For Ohio patients, please only choose one (1) prescription/form.** |               |         |
| CLINICAL INFORMATION  |  |  |               |         |
| <input type="radio"/> B18.2 Chronic Hepatitis C <input type="radio"/> K72.90 <input type="radio"/> K72.91 Hepatic Encephalopathy <input type="radio"/> C22.0 <input type="radio"/> C22.2 <input type="radio"/> C22.7 <input type="radio"/> C22.8 Hepatocellular Carcinoma<br><input type="radio"/> Other _____<br>Genotype: _____ NS5A RAVs <input type="radio"/> Yes <input type="radio"/> No Viral load: _____ IU/ml Viral load date: _____<br><input type="radio"/> Treatment naïve <input type="radio"/> Previously treated: Prior treatment used: _____ <input type="radio"/> Non-responder <input type="radio"/> Responder/Relapser<br>Duration of previous therapy: From _____ to _____ Total of: _____ months HIV Coinfected: <input type="radio"/> Yes <input type="radio"/> No HBV Coinfected: <input type="radio"/> Yes <input type="radio"/> No<br>Compensated Liver Disease: <input type="radio"/> Yes <input type="radio"/> No Cirrhosis: <input type="radio"/> Yes <input type="radio"/> No Metavir Score: _____<br>Solid Organ Transplant recipient: <input type="radio"/> Yes <input type="radio"/> No Awaiting liver transplant: <input type="radio"/> Yes <input type="radio"/> No |  |  |               |         |
| PRESCRIPTION INFORMATION  |  | CHECK TO ENROLL IN PATIENT ASSISTANCE PROGRAM <input type="radio"/>  |               |         |
| MEDICATION  | STRENGTH   | DIRECTIONS   | QTY           | REFILLS |
| <input type="radio"/> Daklinza™<br>(daclatasvir)  | <input type="radio"/> 30mg<br><input type="radio"/> 60mg<br><input type="radio"/> 90mg                                     | Take one tablet PO QD with or without food<br>*Must be taken in combination with sofosbuvir  | 28 day supply |         |
| <input type="radio"/> Epclusa®  | 400mg/100mg tablet   | Take one tablet PO QD with or without food<br><input type="radio"/> Genotype 1, 2, 3, 4, 5, 6 with cirrhosis; 12 wks<br><input type="radio"/> Genotype 1, 2, 3, 4, 5, 6 with decompensated cirrhosis: 12 wks + Ribavirin 12 wks<br>Ribavirin: If <75kg: give 500mg PO twice daily If >75kg: give 600mg PO twice daily  | 28 tablets    |         |
| <input type="radio"/> Harvoni®<br>(ledipasvir/sofosbuvir)   | 90/400mg tablet  | Take one tablet PO QD with or without food<br><input type="radio"/> Naïve without cirrhosis who have a pretreatment HCV RNA <6 million IU/ml; 8 wks<br><input type="radio"/> Naïve with or without cirrhosis; 12 wks<br><input type="radio"/> Experienced without cirrhosis; 12 wks <input type="radio"/> Experienced with cirrhosis; 24 wks   | 28 day supply |         |
| <input type="radio"/> Olysio®<br>(simeprevir)   | 150mg capsule  | Take 150mg (1 capsule) PO QD with food<br>*Monotherapy not recommended   | 28 day supply |         |
| <input type="radio"/> Riba-pak®   | <input type="radio"/> 600mg<br><input type="radio"/> 800mg<br><input type="radio"/> 1000mg<br><input type="radio"/> 1200mg | <input type="radio"/> 200mg every morning, 400mg every evening<br><input type="radio"/> 400mg every morning, 400mg every evening<br><input type="radio"/> 600mg every morning, 400mg every evening<br><input type="radio"/> 600mg every morning, 600mg every evening   | 28 day supply |         |
| <input type="radio"/> Ribasphere®   | 200mg  |  | 28 day supply |         |
| <input type="radio"/> Sovaldi®<br>(sofosbuvir)  | 400mg tablet   | Take 400mg (1 tablet) PO QD with or without food<br>*Monotherapy not recommended   | 28 day supply |         |
| <input type="radio"/> Technivie™<br>(ombitasvir/paritaprevir/ritonavir)   | 12.5/75/50mg   | Take two fixed dose combination tablets daily in the morning<br>*Must be taken with ribavirin  | 28 day supply |         |
| <input type="radio"/> Viekira Pak®<br>(Ombitasvir/Paritaprevir/ritonavir/dasabuvir)   | 12.5/75/50/250 mg  | Take daily as directed<br><input type="radio"/> Genotype 1A without cirrhosis Viekira Pak + ribavirin or Viekira XR + ribavirin; 12 wks*<br><input type="radio"/> Genotype 1A with compensated cirrhosis Viekira Pak + ribavirin or Viekira XR + ribavirin; 24 wks   | 28 day supply |         |
| <input type="radio"/> Viekira XR®<br>(dasabuvir/ombitasvir/paritaprevir/ritonavir)  | 200/8.33/50/33.33mg  | <input type="radio"/> Genotype 1B without cirrhosis Viekira Pak or Viekira XR; 12 wks**<br><input type="radio"/> Genotype 1B with compensated cirrhosis Viekira Pak or Viekira XR; 12 wks**<br>*For liver transplant patients with normal hepatic function and mild fibrosis increase to 24 wks<br>**For liver transplant patients with normal hepatic function and mild fibrosis add ribavirin and increase to 24 wks   |               |         |
| <input type="radio"/> Zepatier™<br>(elbasvir/grazoprevir)   | 50/100mg tablet  | Take one tablet PO QD with or without food; 12 wks<br>Genotype 1a NS5A resistance-associated polymorphisms are present, administer with ribavirin and extend therapy to 16 wks   | 28 day supply |         |
| Date meds needed: _____ <input type="radio"/> New <input type="radio"/> Refill  |  | Ship to: <input type="radio"/> Patient's home <input type="radio"/> Physician's Office <input type="radio"/> Other _____   |               |         |

CONFIDENTIALITY NOTICE: This communication is intended for and should be delivered to the individual or entity to which it is addressed and contains information that is privileged, confidential and exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute or copy this information. Please notify the sender immediately if you have received this document in error.