

Rheumatology Enrollment Form

Ph: 806-324-5447 • Toll Free 866-629-6779
 Toll Free Fax: 866-217-8034
 Email: specialty@maxor.com
www.maxorspecialty.com



PATIENT INFORMATION

Patient Name _____
 Date of Birth _____ Male Female
 Street Address _____ Apt # _____
 City _____ State _____ Zip _____
 Phone-Primary _____ Secondary _____
 English Spanish Other _____
 Email Address _____
 Social Security # _____
 NKDA Known Drug Allergies _____
 Weight _____ kg/lb Height _____ in/cm

Please attach front and back of patient's insurance cards

PHYSICIAN INFORMATION

Physician Name _____
 NPI _____ License # _____
 Office Contact _____
 Street Address _____ Ste # _____
 City _____ State _____ Zip _____
 Phone _____ Fax _____
 By signing this form, I authorize Maxor Specialty to act as my agent for prior authorizations & prescription reimbursement for the above listed patient.
 Physician Signature _____
 Date _____
 Dispense as written Product substitution permitted

** For Ohio patients, please only choose one (1) prescription/form.**

CLINICAL INFORMATION

Diagnosis: M06.9 Rheumatoid Arthritis L40.50 Psoriatic Arthritis M45.9 Ankylosing Spondylitis
 M32.10 Systemic Lupus Erythematosus L40.8 Psoriasis Moderate to Severe Plaque M81.0 Osteoporosis
 Other _____
 Prior (FAILED): Medications _____ Length of Treatment _____ Reason for Discontinuing _____
 Medications _____ Length of Treatment _____ Reason for Discontinuing _____
 TB Test: Yes No Results: _____ Forteo/Prolia: T-Score: _____ Type _____ Date _____

PRESCRIPTION INFORMATION

CHECK TO ENROLL IN PATIENT ASSISTANCE PROGRAM

| MEDICATION | STRENGTH | DIRECTIONS | QTY | REFILLS |
|---------------------------------|---|---|-----|---------|
| <input type="radio"/> Actemra® | <input type="radio"/> 4mg/kg <input type="radio"/> 8mg/kg <input type="radio"/> 162mg/0.9ml PFS | <input type="radio"/> Infuse IV over 1 hr every 4 wks <input type="radio"/> <100kg inject 162mg SQ every other wk, then increase to every wk based on clinical response <input type="radio"/> ≥100kg inject 162kg SQ every wk | | |
| <input type="radio"/> Benlysta® | <input type="radio"/> 10mg/kg | <input type="radio"/> Infuse _____ mg at wks 0, 2, and 4, then every 4 wks thereafter | | |
| <input type="radio"/> Cimzia® | <input type="radio"/> Cimzia Starter Kit <input type="radio"/> 200mg/1ml PFS (2) | Initial Dose: <input type="radio"/> Inject 400mg SQ at wks 0,2,4 Maintenance Dose: <input type="radio"/> Inject 200mg every other wk <input type="radio"/> Inject 400mg SQ every 4 wks | | |
| <input type="radio"/> Cosentyx® | <input type="radio"/> 2 pk (2X150mg)Pen/PFS <input type="radio"/> 150mg (single) <input type="radio"/> Pen <input type="radio"/> PFS | Initial Dose: <input type="radio"/> Inject 150mg SQ at wks 0, 1, 2, 3, 4 <input type="radio"/> Inject 300mg SQ at wks 0, 1, 2, 3, 4 Maintenance Dose: <input type="radio"/> Inject <input type="radio"/> 150mg SQ every 4 wks <input type="radio"/> Inject <input type="radio"/> 300mg SQ every 4 wks | | |
| <input type="radio"/> Enbrel® | <input type="radio"/> 50mg/ml Sureclick <input type="radio"/> 50mg/ml PFS <input type="radio"/> 25mg/0.5ml PFS <input type="radio"/> 25mg Vial | <input type="radio"/> Inject 50mg SQ once a wk <input type="radio"/> Inject 25mg SQ twice a wk (72-96 hrs apart) <input type="radio"/> Other _____ | | |
| <input type="radio"/> Humira® | <input type="radio"/> 40mg <input type="radio"/> Pen <input type="radio"/> PFS | <input type="radio"/> Inject 40mg SQ every other wk <input type="radio"/> Other _____ | | |
| <input type="radio"/> Orencia® | <input type="radio"/> 125mg PFS <input type="radio"/> 250mg Vial | <input type="radio"/> Inject 125mg SQ once a wk <input type="radio"/> >100kg: 1000mg IV wks 0, 2, 4, then Q 4 wks starting wk 8 <input type="radio"/> 60 - 100kg: 750mg IV wks 0, 2, 4, then Q 4 wks starting wk 8 <input type="radio"/> < 60kg: 500mg IV wks 0, 2, 4, then Q 4 wks starting wk 8 | | |
| <input type="radio"/> Otezla® | <input type="radio"/> Starter Pack <input type="radio"/> 30mg Tablet | <input type="radio"/> Take 1 Tablet on day 1 then twice daily as directed <input type="radio"/> Take 1 tablet by mouth twice a day | | |
| <input type="radio"/> Remicade® | <input type="radio"/> 100mg Vial | <input type="radio"/> Infuse _____ mg/kg at wks 0, 2, and 6, then _____ wks thereafter | | |
| <input type="radio"/> Rituxan® | <input type="radio"/> 500mg Vial | <input type="radio"/> Infuse 1000mg IV days 1 and 15, then _____ wks (no sooner than 16 wks) | | |
| <input type="radio"/> Simponi® | <input type="radio"/> 2mg/kg <input type="radio"/> 50mg | <input type="radio"/> Infuse IV over 30 min, repeat dose in 4 wks, then every 8 wks thereafter <input type="radio"/> Inject 50mg SQ monthly | | |
| <input type="radio"/> Stelera® | <input type="radio"/> 45mg <input type="radio"/> 90mg | Initial Dose: <input type="radio"/> 45mg SQ; repeat dose in 4 wks <input type="radio"/> >100kg (220lbs) and with plaque psoriasis 90mg SQ; repeat dose in 4 wks Maintenance Dose: <input type="radio"/> 45mg Q 12 wks starting at 16 wks <input type="radio"/> >100kg(220lbs) and with plaque psoriasis 90mg Q 12 wks starting at 16 wks | | |
| <input type="radio"/> Xeljanz® | <input type="radio"/> 5mg Tablet <input type="radio"/> 11mg XR Tablet | <input type="radio"/> Take 5mg by mouth twice a day <input type="radio"/> Take 11mg by mouth once a day | | |
| <input type="radio"/> Other | | | | |

Date meds needed: _____ New Refill

Ship to: Patient's home Physician's Office Other _____

CONFIDENTIALITY NOTICE: This communication is intended for and should be delivered to the individual or entity to which it is addressed and contains information that is privileged, confidential and exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute or copy this information. Please notify the sender immediately if you have received this document in error.